

Claudin-4 Polyclonal Antibody
Catalog # AP69133**Specification****Claudin-4 Polyclonal Antibody - Product Information**

Application	WB, IHC-P
Primary Accession	O14493
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

Claudin-4 Polyclonal Antibody - Additional Information**Gene ID** 1364**Other Names**

CLDN4; CPER; CPETR1; WBSCR8; Claudin-4; Clostridium perfringens enterotoxin receptor; CPE-R; CPE-receptor; Williams-Beuren syndrome chromosomal region 8 protein

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications.

IHC-P~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

Claudin-4 Polyclonal Antibody - Protein Information**Name** CLDN4 {ECO:0000303|PubMed:35773259, ECO:0000312|HGNC:HGNC:2046}**Function**

Can associate with other claudins to regulate tight junction structural and functional strand dynamics (PubMed:35773259, PubMed:36008380). May coassemble with CLDN8 into tight junction strands containing anion-selective channels that convey paracellular chloride permeability in renal collecting ducts (By similarity) (PubMed:36008380). May integrate into CLDN3 strands to modulate localized tight junction barrier properties (PubMed:35773259, PubMed:36008380). May disrupt strand assembly of channel-forming CLDN2 and CLDN15 and inhibit cation conductance (PubMed:35773259, PubMed:36008380). Cannot form

tight junction strands on its own (PubMed:35773259, PubMed:36008380).

Cellular Location


Cell junction, tight junction {ECO:0000250|UniProtKB:O35054}. Cell membrane; Multi-pass membrane protein

Claudin-4 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Claudin-4 Polyclonal Antibody - Images



Claudin-4 Polyclonal Antibody - Background

Channel-forming tight junction protein that mediates paracellular chloride transport in the kidney. Plays a critical role in the paracellular reabsorption of filtered chloride in the kidney collecting ducts. Claudins play a major role in tight junction-specific obliteration of the intercellular space, through calcium-independent cell-adhesion activity.