

## Lefty (LEFTB) Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP7297b

### **Specification**

## Lefty (LEFTB) Antibody (C-term) - Product Information

**Application** IHC-P, WB,E **Primary Accession** 075610 Reactivity Human **Rabbit** Host Clonality **Polyclonal** Isotype Rabbit IgG Calculated MW 40880 Antigen Region 283-314

### Lefty (LEFTB) Antibody (C-term) - Additional Information

### **Gene ID** 10637

#### **Other Names**

Left-right determination factor 1, Left-right determination factor B, Protein lefty-1, Protein lefty-B, LEFTY1, LEFTB, LEFTYB

### Target/Specificity

This Lefty (LEFTB) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 283-314 amino acids from the C-terminal region of human Lefty (LEFTB).

# **Dilution**

IHC-P~~1:10~50 WB~~1:1000

E~~Use at an assay dependent concentration.

#### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

### **Precautions**

Lefty (LEFTB) Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

# Lefty (LEFTB) Antibody (C-term) - Protein Information

### Name LEFTY1



## Synonyms LEFTB, LEFTYB

Function Required for left-right axis determination as a regulator of LEFTY2 and NODAL.


**Cellular Location** Secreted.

# Lefty (LEFTB) Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

# Lefty (LEFTB) Antibody (C-term) - Images



Western blot analysis of anti-LEFTB(C-term) Pab (Cat.#AP7297b) in HL60 cell line lysates (35ug/lane). LEFTB(arrow) was detected using the purified Pab.





Formalin-fixed and paraffin-embedded human colon carcinoma tissue reacted with LEFTB antibody (C-term) (Cat.#AP7279b), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

## Lefty (LEFTB) Antibody (C-term) - Background

LEFTB is a member of the TGF-beta family of proteins. A similar secreted protein in mouse plays a role in left-right asymmetry determination of organ systems during development. Alternative processing of this protein can yield three different products.

# Lefty (LEFTB) Antibody (C-term) - References

Kosaki K., Am. J. Hum. Genet. 64:712-721(1999) Dvash, T., Stem Cells 25 (2), 465-472 (2007) Besser, D., J. Biol. Chem. 279 (43), 45076-45084 (2004)