

TIF-IA Polyclonal Antibody
Catalog # AP74314**Specification**

TIF-IA Polyclonal Antibody - Product Information

Application	WB
Primary Accession	Q9NYV6
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

TIF-IA Polyclonal Antibody - Additional Information**Gene ID** 54700**Other Names**

RNA polymerase I-specific transcription initiation factor RRN3 (Transcription initiation factor IA) (TIF-IA)

Dilution

WB~~WB 1:500-2000, ELISA 1:10000-20000

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

TIF-IA Polyclonal Antibody - Protein Information**Name** RRN3**Synonyms** TIFIA**Function**

Required for efficient transcription initiation by RNA polymerase I (Pol I). Required for the formation of the competent pre- initiation complex (PIC).

Cellular Location

Nucleus, nucleolus.


TIF-IA Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TIF-IA Polyclonal Antibody - Images



TIF-IA Polyclonal Antibody - Background

Required for efficient transcription initiation by RNA polymerase I. Required for the formation of the competent preinitiation complex (PIC). Dissociates from pol I as a consequence of transcription. In vitro, cannot activate transcription in a subsequent transcription reaction (By similarity).