

Anti-CCL5 / RANTES Reference Antibody (VLST-002)
Recombinant Antibody
Catalog # APR10288**Specification**

Anti-CCL5 / RANTES Reference Antibody (VLST-002) - Product Information

Application	FC, Kinetics, Animal Model
Primary Accession	P13501
Reactivity	Human, Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	150 KDa

Anti-CCL5 / RANTES Reference Antibody (VLST-002) - Additional Information**Target/Specificity**
CCL5 / RANTES**Endotoxin**
< 0.001EU/ µg,determined by LAL method.**Conjugation**
Unconjugated**Expression system**
CHO Cell**Format**
Purified monoclonal antibody supplied in PBS, pH6.0, without preservative.This antibody is purified through a protein A column.**Anti-CCL5 / RANTES Reference Antibody (VLST-002) - Protein Information****Name** CCL5**Synonyms** D17S136E, SCYA5**Function**
Chemoattractant for blood monocytes, memory T-helper cells and eosinophils. Causes the release of histamine from basophils and activates eosinophils. May activate several chemokine receptors including CCR1, CCR3, CCR4 and CCR5. One of the major HIV-suppressive factors produced by CD8+ T-cells. Recombinant RANTES protein induces a dose-dependent inhibition of different strains of HIV-1, HIV-2, and simian immunodeficiency virus (SIV). The processed form RANTES(3-68) acts as a natural chemotaxis inhibitor and is a more potent inhibitor of HIV-1-infection. The second processed form RANTES(4-68) exhibits reduced chemotactic and HIV-suppressive activity compared with RANTES(1-68) and RANTES(3-68) (PubMed:1380064, PubMed:15923218, PubMed:16791620, PubMed:8525373, PubMed:9516414). May also be an agonist of the G protein-coupled receptor GPR75, stimulating inositol trisphosphate production and calcium mobilization through its activation. Together with GPR75, may play a role in neuron survival through activation of a downstream signaling pathway involving the PI3, Akt and MAP kinases. By activating GPR75 may also play a role in insulin secretion by islet cells (PubMed:23979485).

Cellular Location
Secreted.


Tissue Location
Expressed in the follicular fluid (at protein level). T-cell and macrophage specific.

Anti-CCL5 / RANTES Reference Antibody (VLST-002) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-CCL5 / RANTES Reference Antibody (VLST-002) - Images



Anti-CCL5 / RANTES Reference Antibody (VLST-002) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%



The purity of Anti-CCL5 / RANTES Reference Antibody (VLST-002) is more than 95% ,determined by SEC-HPLC.