

**Anti-Integrin  $\alpha$ V / ITGAV / CD51 Reference Antibody (intetumumab)  
Recombinant Antibody  
Catalog # APR10606****Specification****Anti-Integrin  $\alpha$ V / ITGAV / CD51 Reference Antibody (intetumumab) - Product Information**

|                   |                            |
|-------------------|----------------------------|
| Application       | FC, Kinetics, Animal Model |
| Primary Accession | <a href="#">P06756</a>     |
| Reactivity        | Human                      |
| Clonality         | Monoclonal                 |
| Isotype           | IgG1                       |
| Calculated MW     | 145.14 KDa                 |

**Anti-Integrin  $\alpha$ V / ITGAV / CD51 Reference Antibody (intetumumab) - Additional Information****Target/Specificity**Integrin  $\alpha$ V / ITGAV / CD51**Endotoxin**< 0.001EU/  $\mu$ g,determined by LAL method.**Conjugation**

Unconjugated

**Expression system**

CHO Cell

**Format**

Purified monoclonal antibody supplied in PBS, pH6.0, without preservative.This antibody is purified through a protein A column.

**Anti-Integrin  $\alpha$ V / ITGAV / CD51 Reference Antibody (intetumumab) - Protein Information****Name** ITGAV ([HGNC:6150](#))**Function**

The alpha-V (ITGAV) integrins are receptors for vitronectin, cytotactin, fibronectin, fibrinogen, laminin, matrix metalloproteinase- 2, osteopontin, osteomodulin, prothrombin, thrombospondin and vWF. They recognize the sequence R-G-D in a wide array of ligands. ITGAV:ITGB3 binds to fractalkine (CX3CL1) and may act as its coreceptor in CX3CR1- dependent fractalkine signaling (PubMed:<a href="http://www.uniprot.org/citations/23125415" target="\_blank">23125415</a>). ITGAV:ITGB3 binds to NRG1 (via EGF domain) and this binding is essential for NRG1-ERBB signaling (PubMed:<a href="http://www.uniprot.org/citations/20682778" target="\_blank">20682778</a>). ITGAV:ITGB3 binds to FGF1 and this binding is essential for FGF1 signaling (PubMed:<a href="http://www.uniprot.org/citations/18441324" target="\_blank">18441324</a>). ITGAV:ITGB3 binds to FGF2 and this binding is essential for FGF2 signaling (PubMed:<a href="http://www.uniprot.org/citations/28302677" target="\_blank">28302677</a>).

target="\_blank">28302677</a>). ITGAV:ITGB3 binds to IGF1 and this binding is essential for IGF1 signaling (PubMed:<a href="http://www.uniprot.org/citations/19578119" target="\_blank">19578119</a>). ITGAV:ITGB3 binds to IGF2 and this binding is essential for IGF2 signaling (PubMed:<a href="http://www.uniprot.org/citations/28873464" target="\_blank">28873464</a>). ITGAV:ITGB3 binds to IL1B and this binding is essential for IL1B signaling (PubMed:<a href="http://www.uniprot.org/citations/29030430" target="\_blank">29030430</a>). ITGAV:ITGB3 binds to PLA2G2A via a site (site 2) which is distinct from the classical ligand-binding site (site 1) and this induces integrin conformational changes and enhanced ligand binding to site 1 (PubMed:<a href="http://www.uniprot.org/citations/18635536" target="\_blank">18635536</a>, PubMed:<a href="http://www.uniprot.org/citations/25398877" target="\_blank">25398877</a>). ITGAV:ITGB3 and ITGAV:ITGB6 act as receptors for fibrillin-1 (FBN1) and mediate R-G-D-dependent cell adhesion to FBN1 (PubMed:<a href="http://www.uniprot.org/citations/12807887" target="\_blank">12807887</a>, PubMed:<a href="http://www.uniprot.org/citations/17158881" target="\_blank">17158881</a>). Integrin alpha-V/beta-6 or alpha-V/beta-8 (ITGAV:ITGB6 or ITGAV:ITGB8) mediates R-G-D-dependent release of transforming growth factor beta-1 (TGF-beta-1) from regulatory Latency-associated peptide (LAP), thereby playing a key role in TGF-beta-1 activation (PubMed:<a href="http://www.uniprot.org/citations/15184403" target="\_blank">15184403</a>, PubMed:<a href="http://www.uniprot.org/citations/22278742" target="\_blank">22278742</a>, PubMed:<a href="http://www.uniprot.org/citations/28117447" target="\_blank">28117447</a>). ITGAV:ITGB3 acts as a receptor for CD40LG (PubMed:<a href="http://www.uniprot.org/citations/31331973" target="\_blank">31331973</a>). ITGAV:ITGB3 acts as a receptor for IBSP and promotes cell adhesion and migration to IBSP (PubMed:<a href="http://www.uniprot.org/citations/10640428" target="\_blank">10640428</a>).

#### **Cellular Location**


Cell membrane; Single-pass type I membrane protein. Cell junction, focal adhesion


#### **Anti-Integrin aV / ITGAV / CD51 Reference Antibody (intetumumab) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **Anti-Integrin aV / ITGAV / CD51 Reference Antibody (intetumumab) - Images**





Anti-Integrin  $\alpha$ V / ITGAV / CD51 Reference Antibody (intetumumab) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-Integrin  $\alpha$ V / ITGAV / CD51 Reference Antibody (intetumumab) is more than 95%, determined by SEC-HPLC.