

BAD Antibody
Rabbit Anti-Human BAD Polyclonal
Catalog # ASM10622**Specification**

BAD Antibody - Product Information

Application	IHC
Primary Accession	O92934
Other Accession	NP_004313.1
Host	Rabbit
Clonality	Polyclonal
Format	FITC
Target/Specificity	
BAD	

Other Names

Bcl2 antagonist of cell death Antibody, BAD Antibody, Bcl2-L-8 Antibody, Bcl-xL/Bcl-2-associated death promoter Antibody, Bcl2-associated agonist of cell death Antibody, Bcl-2-binding component 6 Antibody, BBC6 Antibody, BCL2L8 Antibody, Bcl-2-like protein 8 Antibody

Immunogen

Synthetic peptide from the mid-protein of Human BAD (aa. 60-90)

Purification

Peptide Affinity Purified

Storage -20°C

Storage Buffer

PBS pH 7.4, 50% glycerol, 0.09% sodium azide *Storage buffer may change when conjugated

Shipping Temperature Blue Ice or 4°C


Cellular Localization


Mitochondrion Outer Membrane | Cytoplasm

BAD Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BAD Antibody - Images



Immunohistochemistry analysis using Rabbit Anti-BAD Polyclonal Antibody (SPC-773). Tissue: Prostate. Species: Human. Fixation: Formalin fixed paraffin-embedded. Primary Antibody: Rabbit Anti-BAD Polyclonal Antibody (SPC-773). | Western blot analysis of Rat Kidney showing detection of ~18.3 kDa BAD protein using Rabbit Anti-BAD Polyclonal Antibody (SPC-773). Lane 1: Molecular Weight Ladder (MW). Lane 2: Rat Kidney. Load: 15 μ g. Block: 5% Skim Milk in 1X TBST. Primary Antibody: Rabbit Anti-BAD Polyclonal Antibody (SPC-773) at 1:1000 for 2 hours at RT. Secondary Antibody: Goat Anti-Rabbit IgG: HRP at 1:3000 for 1 hour at RT. Color Development: ECL solution for 5 min at RT. Predicted/Observed Size: ~18.3 kDa. Other Band(s): 28 kDa (PTM).



Immunohistochemistry analysis using Rabbit Anti-BAD Polyclonal Antibody (SPC-773). Tissue: Prostate. Species: Human. Fixation: Formalin fixed paraffin-embedded. Primary Antibody: Rabbit Anti-BAD Polyclonal Antibody (SPC-773). | Western blot analysis of Rat Kidney showing detection of ~18.3 kDa BAD protein using Rabbit Anti-BAD Polyclonal Antibody (SPC-773). Lane 1: Molecular Weight Ladder (MW). Lane 2: Rat Kidney. Load: 15 μ g. Block: 5% Skim Milk in 1X TBST. Primary Antibody: Rabbit Anti-BAD Polyclonal Antibody (SPC-773) at 1:1000 for 2 hours at RT. Secondary Antibody: Goat Anti-Rabbit IgG: HRP at 1:3000 for 1 hour at RT. Color Development: ECL solution for 5 min at RT. Predicted/Observed Size: ~18.3 kDa. Other Band(s): 28 kDa (PTM).