

A0573

Leader in Biomolecular Solutions for Life Science



EIF3A Rabbit pAb

Catalog No.: A0573

4 Publications

Basic Information

Observed MW

166kDa

Calculated MW

167kDa

Category

Polyclonal Antibody

Applications

WB,IF/ICC,ELISA

Cross-Reactivity

Human,Mouse

Background

Enables RNA binding activity. Contributes to translation initiation factor activity. Involved in IRES-dependent viral translational initiation; formation of cytoplasmic translation initiation complex; and viral translational termination-reinitiation. Located in cytosol; nucleolus; and nucleoplasm. Part of eukaryotic translation initiation factor 3 complex.

Recommended Dilutions

WB 1:1000 - 1:5000

IF/ICC 1:50 - 1:100

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

8661

Swiss Prot

Q14152

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

EIF3; P167; p180; p185; TIF32; EIF3S10; eIF3-p170; eIF3-theta; EIF3A

Contact

 www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of lysates from RAW 264.7 cells, using EIF3A Rabbit pAb (A0573) at 1:2000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.



Immunofluorescence analysis of U2OS cells using EIF3A Rabbit pAb (A0573) at dilution of 1:100.

Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.