# ABclonal®

# **DDR2 Rabbit pAb**

Catalog No.: A10060 1 Publications

#### **Basic Information**

#### **Observed MW**

110kDa

#### **Calculated MW**

97kDa

#### Category

Polyclonal Antibody

#### **Applications**

WB,ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

This gene encodes a member of the discoidin domain receptor subclass of the receptor tyrosine kinase (RTKs) protein family. RTKs play a key role in the communication of cells with their microenvironment. The encoded protein is a collagen-induced receptor that activates signal transduction pathways involved in cell adhesion, proliferation, and extracellular matrix remodeling. This protein is expressed in numerous cell types and may alos be involved in wound repair and regulate tumor growth and invasiveness. Mutations in this gene are the cause of short limb-hand type spondylometaepiphyseal dysplasia.

#### **Recommended Dilutions**

WB

1:500 - 1:2000

# **Immunogen Information**

Gene ID 4921 **Swiss Prot** 

Q16832

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 290-400 of human DDR2 (NP\_001014796.1).

#### **Synonyms**

TKT; WRCN; MIG20a; NTRKR3; TYRO10; DDR2

## Contact

•

www.abclonal.com

#### **Product Information**

Source

Isotype

**Purification** 

Rabbit

IgG


Affinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# Validation Data



Western blot analysis of extracts of various cell lines, using DDR2 antibody (A10060) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.