# ABclonal®

## CD53 Rabbit pAb

Catalog No.: A10079

#### **Basic Information**

#### **Observed MW**

35-45kDa

#### **Calculated MW**

24kDa

#### Category

Polyclonal Antibody

#### **Applications**

WB,ELISA

#### **Cross-Reactivity**

Human, Mouse

## **Background**

The protein encoded by this gene is a member of the transmembrane 4 superfamily, also known as the tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. This encoded protein is a cell surface glycoprotein that is known to complex with integrins. It contributes to the transduction of CD2-generated signals in T cells and natural killer cells and has been suggested to play a role in growth regulation. Familial deficiency of this gene has been linked to an immunodeficiency associated with recurrent infectious diseases caused by bacteria, fungi and viruses. Alternative splicing results in multiple transcript variants.

## **Recommended Dilutions**

WB

1:1000 - 1:4000

## **Immunogen Information**

**Gene ID** 

**Swiss Prot** 

963

P19397

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 100-180 of human CD53 (NP\_000551.1).

## **Synonyms**

MOX44; TSPAN25; CD53

## Contact

€

www.abclonal.com

#### **Product Information**

**Source** Rabbit Isotype

IgG

Purification


Affinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

## Validation Data



Western blot analysis of Mouse spleen, using CD53 Rabbit pAb (A10079) at 1:400 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25ug per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.