# ABclonal®

## **RAB34 Rabbit pAb**

Catalog No.: A10332

## **Basic Information**

#### **Observed MW**

29kDa

#### **Calculated MW**

21kDa/28kDa/29kDa

#### Category

Polyclonal Antibody

#### **Applications**

WB, ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

This gene encodes a protein belonging to the RAB family of proteins, which are small GTPases involved in protein transport. This family member is a Golgi-bound member of the secretory pathway that is involved in the repositioning of lysosomes and the activation of macropinocytosis. Alternative splicing of this gene results in multiple transcript variants. An alternatively spliced transcript variant produces the nine-amino acid residue-repeats (NARR) protein, which is a functionally distinct nucleolar protein resulting from a different reading frame.

## **Recommended Dilutions**

WB 1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID**83871

Swiss Prot
PODI83/Q9BZG1

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

## **Synonyms**

RAH; NARR; RAB39; RAB34

## **Contact**

www.abclonal.com

## **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

## **Validation Data**



Western blot analysis of various lysates using RAB34 Rabbit pAb (A10332) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.