

A10389

Leader in Biomolecular Solutions for Life Science



RAB27B Rabbit pAb

Catalog No.: A10389

4 Publications

Basic Information

Observed MW

27kDa

Calculated MW

25kDa

Category

Polyclonal Antibody

Applications

WB, ELISA

Cross-Reactivity

Human, Mouse

Background

Enables guanyl ribonucleotide binding activity; myosin V binding activity; and protein domain specific binding activity. Involved in multivesicular body sorting pathway and positive regulation of exocytosis. Located in Golgi stack and cytoplasmic vesicle.

Recommended Dilutions

WB 1:500 - 1:1000

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

5874

Swiss Prot

O00194

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

C25KG; RAB27B

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of various lysates, using RAB27B Rabbit pAb (A10389) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 25s.



Western blot analysis of lysates from Mouse stomach, using RAB27B Rabbit pAb (A10389) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.