

A11223

Leader in Biomolecular Solutions for Life Science



## Ku70 Rabbit mAb

Catalog No.: A11223

Recombinant

1 Publications

### Basic Information

#### Observed MW

70kDa

#### Calculated MW

70kDa

#### Category

SMab Recombinant Monoclonal  
Antibody

#### Applications

WB,IHC-P,IF/ICC,ELISA

#### Cross-Reactivity

Human,Mouse,Rat

#### CloneNo number

ARC0551

### Recommended Dilutions

**WB** 1:3000 - 1:20000

**IHC-P** 1:200 - 1:800

**IF/ICC** 1:100 - 1:500

**ELISA** Recommended starting  
concentration is 1  
µg/mL. Please optimize  
the concentration  
based on your specific  
assay requirements.

### Contact



[www.abclonal.com](http://www.abclonal.com)

### Background

The p70/p80 autoantigen is a nuclear complex consisting of two subunits with molecular masses of approximately 70 and 80 kDa. The complex functions as a single-stranded DNA-dependent ATP-dependent helicase. The complex may be involved in the repair of nonhomologous DNA ends such as that required for double-strand break repair, transposition, and V(D)J recombination. High levels of autoantibodies to p70 and p80 have been found in some patients with systemic lupus erythematosus.

### Immunogen Information

#### Gene ID

2547

#### Swiss Prot

P12956

#### Immunogen

A synthetic peptide corresponding to a sequence within amino acids 500-609 of human Ku70 (P12956).

#### Synonyms

ML8; KU70; TLAA; CTC75; CTCBF; G22P1; Ku70

### Product Information

#### Source

Rabbit

#### Isotype

IgG

#### Purification


Affinity purification

#### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,0.05% BSA,50% glycerol,pH7.3.


## Validation Data




Western blot analysis of lysates from A549 cells using Ku70 Rabbit mAb (A11223) at 1:6000 dilution incubated overnight at 4°C.  
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
 Lysates/proteins: 25 µg per lane.  
 Blocking buffer: 3% nonfat dry milk in TBST.  
 Detection: ECL Basic Kit (RM00020).  
 Exposure time: 10s.




Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma tissue using Ku70 Rabbit mAb (A11223) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human esophagus tissue using Ku70 Rabbit mAb (A11223) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human tonsil tissue using Ku70 Rabbit mAb (A11223) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Confocal imaging of HeLa cells using Ku70 Rabbit mAb (A11223, dilution 1:100) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG



Confocal imaging of NIH/3T3 cells using Ku70 Rabbit mAb (A11223, dilution 1:100) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG



Confocal imaging of C6 cells using Ku70 Rabbit mAb (A11223, dilution 1:100) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG

Validation Data

---

(H+L) Ab (AS076, dilution 1:500)  
(Green). DAPI was used for nuclear  
staining (Blue). Objective: 100x.

(H+L) Ab (AS076, dilution 1:500)  
(Green). DAPI was used for nuclear  
staining (Blue). Objective: 100x.

(H+L) Ab (AS076, dilution 1:500)  
(Green). DAPI was used for nuclear  
staining (Blue). Objective: 100x.