

A11412

Leader in Biomolecular Solutions for Life Science



# Histone H2AX Rabbit mAb

Catalog No.: A11412

Recombinant

4 Publications

## Basic Information

### Observed MW

Refer to figures

### Calculated MW

15kDa

### Category

SMab Recombinant Monoclonal  
Antibody

### Applications

IHC-P,ChIP,ELISA

### Cross-Reactivity

Human,Mouse,Rat,Other (Wide Range  
Predicted)

### CloneNo number

ARC0590

## Recommended Dilutions

IHC-P 1:1000 - 1:4000

ChIP 5µg antibody for  
5µg-10µg of Chromatin

ELISA Recommended starting  
concentration is 1  
µg/mL. Please optimize  
the concentration  
based on your specific  
assay requirements.

## Contact



[www.abclonal.com](http://www.abclonal.com)

## Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene encodes a replication-independent histone that is a member of the histone H2A family, and generates two transcripts through the use of the conserved stem-loop termination motif, and the polyA addition motif.

## Immunogen Information

Gene ID  
3014

Swiss Prot  
P16104

### Immunogen

A synthetic peptide corresponding to a sequence within amino acids 1-100 of human Histone H2AX (P16104).

### Synonyms

H2A.X; H2A/X; H2AFX; Histone H2AX

## Product Information

Source  
Rabbit


Isotype  
IgG

Purification  
Affinity purification


### Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS with 0.02% sodium azide,0.05% BSA,50% glycerol,pH7.3.




Immunohistochemistry analysis of paraffin-embedded Mouse brain tissue using Histone H2AX Rabbit mAb (A11412) at a dilution of 1:4000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat brain tissue using Histone H2AX Rabbit mAb (A11412) at a dilution of 1:4000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat testis tissue using Histone H2AX Rabbit mAb (A11412) at a dilution of 1:4000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Chromatin immunoprecipitation analysis of extracts of HeLa cells, using Histone H2AX Rabbit mAb antibody (A11412) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.