# ABclonal www.abclonal.com

## MC3R Rabbit pAb

Catalog No.: A11609

## **Basic Information**

## **Observed MW**

36kDa

#### **Calculated MW**

36kDa

#### **Category**

Polyclonal Antibody

#### **Applications**

WB,ELISA

#### **Cross-Reactivity**

Mouse,Rat

## **Background**

This gene encodes a G-protein-coupled receptor for melanocyte-stimulating hormone and adrenocorticotropic hormone that is expressed in tissues other than the adrenal cortex and melanocytes. This gene maps to the same region as the locus for benign neonatal epilepsy. Mice deficient for this gene have increased fat mass despite decreased food intake, suggesting a role for this gene product in the regulation of energy homeostasis. Mutations in this gene are associated with a susceptibility to obesity in humans.

## **Recommended Dilutions**

**WB** 1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID**Swiss Prot
4159
P41968

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-120 of human MC3R (NP\_063941.3).

## **Synonyms**

MC3; OB20; OQTL; BMIQ9; MC3-R; MC3R

## Contact

www.abclonal.com


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

## Validation Data



Western blot analysis of various lysates using MC3R Rabbit pAb (A11609) at 1:3000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000  $\,$ 

dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.