# ABclonal www.abclonal.com

## Hsp90α Rabbit pAb

Catalog No.: A12448 4 Publications

## **Basic Information**

#### **Observed MW**

101kDa

#### **Calculated MW**

85kDa

## Category

Polyclonal Antibody

#### **Applications**

WB,IHC-P,IF/ICC,ELISA

## **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

The protein encoded by this gene is an inducible molecular chaperone that functions as a homodimer. The encoded protein aids in the proper folding of specific target proteins by use of an ATPase activity that is modulated by co-chaperones. Two transcript variants encoding different isoforms have been found for this gene.

## **Recommended Dilutions**

WB	1:500 - 1:2000
IHC-P	1:50 - 1:200
IF/ICC	1:50 - 1:200

ELISA Recommended starting concentration is 1 

µg/mL. Please optimize 
the concentration

based on your specific assay requirements.

## **Contact**

•	www.abclonal.com
	www.abcionai.com

## **Immunogen Information**

Gene ID	Swiss Prot	
3320	P07900	

## **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 629-732 of human  $Hsp90\alpha$  (NP\_005339.3).

## **Synonyms**


EL52; HSPN; LAP2; HSP86; HSPC1; HSPCA; Hsp89; Hsp90; LAP-2; HSP89A; HSP90A; HSP90N; Hsp103; HSPCAL1; HSPCAL4; HEL-S-65p; Hsp90 $\alpha$ 

## **Product Information**

Source	Isotype	Purification
Rabbit	IgG	Affinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.09% Sodium azide,50% glycerol,pH7.3.



Western blot analysis of various lysates using Hsp90α Rabbit pAb (A12448) at 1:1000

dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000


dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 30s.




Immunohistochemistry analysis of paraffin-embedded Rat testis using Hsp90α Rabbit pAb (A12448) at dilution of 1:200 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse brain using Hsp90α Rabbit pAb (A12448) at dilution of 1:200 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat brain using Hsp90α Rabbit pAb (A12448) at dilution of 1:200 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunofluorescence analysis of C6 cells using Hsp90α Rabbit pAb (A12448) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of paraffin-embedded Mouse testis using Hsp90α Rabbit pAb (A12448) at dilution of 1:200 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.