# **NUP35 Rabbit pAb**

Catalog No.: A12762



### **Basic Information**

#### **Observed MW**

37kDa

### **Calculated MW**

35kDa

### Category

Polyclonal Antibody

### **Applications**

WB, ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

This gene encodes a member of the nucleoporin family. The encoded protein contains two membrane binding regions, is localized to the nuclear rim, and is part of the nuclear pore complex. All molecules entering or leaving the nucleus either diffuse through or are actively transported by the nuclear pore complex. Alternative splicing results in multiple transcript variants. Pseudogenes of this gene have been defined on chromosomes 7 and 10.

# **Recommended Dilutions**

WB 1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID**Swiss Prot
129401
Q8NFH5

### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

# **Synonyms**

MP44; NP44; MP-44; NUP53; NUP35

### **Contact**

www.abclonal.com


# **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

# Validation Data



Western blot analysis of various lysates using NUP35 Rabbit pAb (A12762) at 1:3000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 5s.