# **IGFBP7 Rabbit pAb**

Catalog No.: A13023



# **Basic Information**

#### **Observed MW**

35kDa

### **Calculated MW**

29kDa

### **Category**

Polyclonal Antibody

### **Applications**

WB,ELISA

### **Cross-Reactivity**

Mouse,Rat

## **Background**

This gene encodes a member of the insulin-like growth factor (IGF)-binding protein (IGFBP) family. IGFBPs bind IGFs with high affinity, and regulate IGF availability in body fluids and tissues and modulate IGF binding to its receptors. This protein binds IGF-I and IGF-II with relatively low affinity, and belongs to a subfamily of low-affinity IGFBPs. It also stimulates prostacyclin production and cell adhesion. Alternatively spliced transcript variants encoding different isoforms have been described for this gene, and one variant has been associated with retinal arterial macroaneurysm (PMID:21835307).

# **Recommended Dilutions**

WB 1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

### **Immunogen Information**

**Gene ID**Swiss Prot
Q16270
Q16270

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 27-282 of human IGFBP7 (NP\_001544.1).

### **Synonyms**

AGM; PSF; TAF; FSTL2; IBP-7; MAC25; IGFBP-7; RAMSVPS; IGFBP-7v; IGFBPRP1; IGFBP7

### Contact

www.abclonal.com


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

# Validation Data



Western blot analysis of various lysates using IGFBP7 Rabbit pAb (A13023) at 1:1000

dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000  $\,$ 

dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.