# ABclonal® www.abclonal.com

## **GLYAT Rabbit pAb**

Catalog No.: A13831

### **Basic Information**

#### **Observed MW**

36kDa

#### **Calculated MW**

34kDa

#### **Category**

Polyclonal Antibody

## **Applications**

WB, ELISA

#### **Cross-Reactivity**

Human

## **Background**

The glycine-N-acyltransferase protein conjugates glycine with acyl-CoA substrates in the mitochondria. The protein is thought to be important in the detoxification of endogenous and xenobiotic acyl-CoA's. Two transcript variants encoding different isoforms have been found for this gene.

## **Recommended Dilutions**

**WB** 1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID**Swiss Prot
10249
Q6IB77

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

## **Synonyms**

CAT; GAT; ACGNAT; GLYAT

## **Contact**

www.abclonal.com


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

## Validation Data



Western blot analysis of lysates from 293T cells, using GLYAT Rabbit pAb (A13831) at 1:3000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.