# ABclonal® www.abclonal.com

## **IL31 Rabbit pAb**

Catalog No.: A14982

## **Basic Information**

#### **Observed MW**

18kDa

#### **Calculated MW**

18kDa

#### Category

Polyclonal Antibody

## **Applications**

WB, ELISA

## **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

IL31, which is made principally by activated Th2-type T cells, interacts with a heterodimeric receptor consisting of IL31RA (MIM 609510) and OSMR (MIM 601743) that is constitutively expressed on epithelial cells and keratinocytes. IL31 may be involved in the promotion of allergic skin disorders and in regulating other allergic diseases, such as asthma (Dillon et al., 2004 [PubMed 15184896]).

## **Recommended Dilutions**

WB 1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID**386653 **Swiss Prot**Q6EBC2

### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

#### **Synonyms**

IL-31; IL31

## **Contact**

www.abclonal.com


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

## Validation Data



Western blot analysis of various lysates using IL31 Rabbit pAb (A14982) at 1:1000

dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000  $\,$ 

dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.