

A15017

Leader in Biomolecular Solutions for Life Science



AARS Rabbit pAb

Catalog No.: A15017

2 Publications

Basic Information

Observed MW

107kDa

Calculated MW

107kDa

Category

Polyclonal Antibody

Applications

WB,IHC-P,IF/ICC,ELISA

Cross-Reactivity

Human,Mouse,Rat

Background

The human alanyl-tRNA synthetase (AARS) belongs to a family of tRNA synthetases, of the class II enzymes. Class II tRNA synthetases evolved early in evolution and are highly conserved. This is reflected by the fact that 498 of the 968-residue polypeptide human AARS shares 41% identity with the E.coli protein. tRNA synthetases are the enzymes that interpret the RNA code and attach specific amino acids to the tRNAs that contain the cognate trinucleotide anticodons. They consist of a catalytic domain which interacts with the amino acid acceptor-T ψ C helix of the tRNA, and a second domain which interacts with the rest of the tRNA structure.

Recommended Dilutions

WB 1:500 - 1:2000

IHC-P 1:50 - 1:200

IF/ICC 1:50 - 1:200

ELISA Recommended starting concentration is 1 μ g/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

16

Swiss Prot

P49588

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

AARS; TTD8; CMT2N; DEE29; HDLS2; EIEE29

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH 7.3.

Contact



www.abclonal.com

Validation Data



Western blot analysis of various lysates using AARS Rabbit pAb (A15017) at 1:3000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 10s.




Immunohistochemistry analysis of paraffin-embedded Rat spleen using AARS Rabbit pAb (A15017) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human lung cancer using AARS Rabbit pAb (A15017) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse liver using AARS Rabbit pAb (A15017) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunofluorescence analysis of 293T cells using AARS Rabbit pAb (A15017) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.