

A15269

Leader in Biomolecular Solutions for Life Science



CTNNA2 Rabbit pAb

Catalog No.: A15269

1 Publications

Basic Information

Observed MW

102kDa

Calculated MW

105kDa

Category

Polyclonal Antibody

Applications

WB,IHC-P,IF/ICC,ELISA

Cross-Reactivity

Human,Mouse,Rat

Background

Enables actin filament binding activity. Involved in negative regulation of Arp2/3 complex-mediated actin nucleation; regulation of neuron migration; and regulation of neuron projection development. Located in cytoplasm. Implicated in complex cortical dysplasia with other brain malformations.

Recommended Dilutions

WB	1:500 - 1:1000
IHC-P	1:50 - 1:200
IF/ICC	1:50 - 1:200
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Contact



www.abclonal.com

Immunogen Information

Gene ID

1496

Swiss Prot

P26232

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

CAPR; CTNR; CAP-R; CT114; CDCBM9; CTNNA2

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of various lysates using CTNNA2 Rabbit pAb (A15269) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 5s.




Immunohistochemistry analysis of paraffin-embedded Rat liver using CTNNA2 Rabbit pAb (A15269) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse liver using CTNNA2 Rabbit pAb (A15269) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse spinal cord using CTNNA2 Rabbit pAb (A15269) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunofluorescence analysis of Neuro-2a cells using CTNNA2 Rabbit pAb (A15269) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of PC-12 cells using CTNNA2 Rabbit pAb (A15269) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.