

A15611

Leader in Biomolecular Solutions for Life Science



GPR109A/HM74A/HCAR2 Rabbit pAb

Catalog No.: A15611

5 Publications

Basic Information

Observed MW

42kDa

Calculated MW

42kDa

Category

Polyclonal Antibody

Applications

WB,IHC-P,IF/ICC,ELISA

Cross-Reactivity

Human,Mouse,Rat

Background

Predicted to enable nicotinic acid receptor activity. Involved in neutrophil apoptotic process and positive regulation of neutrophil apoptotic process. Located in cell junction and plasma membrane.

Recommended Dilutions

WB 1:500 - 1:1000

IHC-P 1:50 - 1:200

IF/ICC 1:50 - 1:200

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

338442

Swiss Prot

Q8TDS4

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

HCA2; HM74a; HM74b; PUMAG; NIACR1; Puma-g; GPR109A; GPR109A/HM74A/HCAR2

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS with 0.09% Sodium azide,50% glycerol,pH7.3.

Contact




www.abclonal.com


Validation Data




Western blot analysis of lysates from BxPC-3 cells using GPR109A/HM74A/HCAR2 Rabbit pAb (A15611) at 1:400 dilution.
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25 µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 180s.




Immunohistochemistry analysis of paraffin-embedded Human colon tissue using GPR109A/HM74A/HCAR2 Rabbit pAb (A15611) at a dilution of 1:100 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human spleen tissue using GPR109A/HM74A/HCAR2 Rabbit pAb (A15611) at a dilution of 1:100 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunofluorescence analysis of mouse spleen cells using GPR109A/HM74A/HCAR2 Rabbit pAb (A15611) at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IF staining protocol. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of rat spleen cells using GPR109A/HM74A/HCAR2 Rabbit pAb (A15611) at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer

Validation Data

pH 6.0 before commencing with IF staining protocol. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.