# ABclonal www.abclonal.com

## Sin3A Rabbit pAb

Catalog No.: A1577 3 Publications

## **Basic Information**

## **Observed MW**

160kDa

#### **Calculated MW**

145kDa

#### Category

Polyclonal Antibody

## **Applications**

WB,IF/ICC,IP,ELISA

## **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

The protein encoded by this gene is a transcriptional regulatory protein. It contains paired amphipathic helix (PAH) domains, which are important for protein-protein interactions and may mediate repression by the Mad-Max complex.

## **Recommended Dilutions**

WB 1:100 - 1:500

**IF/ICC** 1:50 - 1:200

**IP** 0.5μg-4μg antibody for

200µg-400µg extracts

of whole cells

**ELISA** Recommended starting

concentration is 1 µg/mL. Please optimize the concentration based on your specific

assay requirements.

## **Contact**

www.abclonal.com

## **Immunogen Information**

**Gene ID**25942

Swiss Prot
Q96ST3

## **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

## **Synonyms**

WITKOS; DEL15Q24; CHR15DELq24; Sin3A


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.



Western blot analysis of various lysates using Sin3A Rabbit pAb (A1577) at 1:500

dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000


dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.



Immunofluorescence analysis of U2OS cells using Sin3A Rabbit pAb (A1577). Secondary antibody: Cy3conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunoprecipitation analysis of 300  $\mu g$  extracts of Jurkat cells using 3  $\mu g$  Sin3A antibody (A1577). Western blot was performed from the immunoprecipitate using Sin3A antibody (A1577) at a dilution of 1:1000.