ABclonal www.abclonal.com

PTPN5 Rabbit pAb

Catalog No.: A16585

Basic Information

Observed MW

46kDa/61kDa

Calculated MW

64kDa

Category

Polyclonal Antibody

Applications

WB,IF/ICC,ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

Enables phosphotyrosine residue binding activity. Predicted to be involved in peptidyltyrosine dephosphorylation. Predicted to act upstream of or within protein dephosphorylation. Predicted to be located in nucleoplasm. Predicted to be integral component of membrane. Biomarker of Alzheimer's disease.

Recommended Dilutions

WB 1:500 - 1:1000

IF/ICC 1:50 - 1:200

ELISA Recommended starting

concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene IDSwiss Prot
84867
P54829

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

STEP; STEP61; PTPSTEP; PTPN5

Contact


www.abclonal.com

Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.



Western blot analysis of various lysates using PTPN5 Rabbit pAb (A16585) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000


dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.



Immunofluorescence analysis of Neuro-2a cells using PTPN5 Rabbit pAb (A16585) at dilution of 1:100 (40x lens). Secondary antibody: Cy3conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of SH-SY5Y cells using PTPN5 Rabbit pAb (A16585) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.