# TNFSF11 Rabbit pAb

Catalog No.: A17025



### **Basic Information**

#### **Observed MW**

Refer to figures

#### **Calculated MW**

35kDa

#### Category

Polyclonal Antibody

### **Applications**

WB,IHC-P,ELISA

### **Cross-Reactivity**

Human, Mouse

## **Background**

This gene encodes a member of the tumor necrosis factor (TNF) cytokine family which is a ligand for osteoprotegerin and functions as a key factor for osteoclast differentiation and activation. This protein was shown to be a dentritic cell survival factor and is involved in the regulation of T cell-dependent immune response. T cell activation was reported to induce expression of this gene and lead to an increase of osteoclastogenesis and bone loss. This protein was shown to activate antiapoptotic kinase AKT/PKB through a signaling complex involving SRC kinase and tumor necrosis factor receptor-associated factor (TRAF) 6, which indicated this protein may have a role in the regulation of cell apoptosis. Targeted disruption of the related gene in mice led to severe osteopetrosis and a lack of osteoclasts. The deficient mice exhibited defects in early differentiation of T and B lymphocytes, and failed to form lobulo-alveolar mammary structures during pregnancy. Two alternatively spliced transcript variants have been found.

# **Recommended Dilutions**

**WB** 1:500 - 1:2000

**IHC-P** 1:50 - 1:200

**ELISA** Recommended starting

concentration is 1 µg/mL. Please optimize the concentration based on your specific

ased on your specific assay requirements.

# **Immunogen Information**

**Gene ID**Swiss Prot
8600
O14788

### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

## **Synonyms**

ODF; OPGL; sOdf; CD254; OPTB2; RANKL; TNLG6B; TRANCE; hRANKL2; TNFSF11

### **Contact**

www.abclonal.com

### **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

# Validation Data



Immunohistochemistry analysis of paraffin-embedded Human tonsil using TNFSF11 Rabbit pAb (A17025) at dilution of 1:200 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human placenta using TNFSF11 Rabbit pAb (A17025) at dilution of 1:200 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.