

A17165

Leader in Biomolecular Solutions for Life Science



UGT1A6 Rabbit pAb

Catalog No.: A17165

Basic Information

Observed MW

75kDa

Calculated MW

61kDa

Category

Antibody

Applications

WB,ELISA

Cross-Reactivity

Human,Mouse

Background

This gene encodes a UDP-glucuronosyltransferase, an enzyme of the glucuronidation pathway that transforms small lipophilic molecules, such as steroids, bilirubin, hormones, and drugs, into water-soluble, excretable metabolites. This gene is part of a complex locus that encodes several UDP-glucuronosyltransferases. The locus includes thirteen unique alternate first exons followed by four common exons. Four of the alternate first exons are considered pseudogenes. Each of the remaining nine 5' exons may be spliced to the four common exons, resulting in nine proteins with different N-termini and identical C-termini. Each first exon encodes the substrate binding site, and is regulated by its own promoter. The enzyme encoded by this gene is active on phenolic and planar compounds. Alternative splicing in the unique 5' end of this gene results in two transcript variants.

Recommended Dilutions

WB 1:500 - 1:1000

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

54578

Swiss Prot

P19224

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

GNT1; UGT1; HLUGP; UDPGT; UGT1A; UGT1C; UGT1E; UGT1F; HLUGP1; UGT-1A; UGT-1C; UGT-1E; UGT-1F; UGT1.1; UGT1.3; UGT1.5; UGT1.6; UGT1A1; UGT1A3; UGT1A5; UGT1-01; UGT1-03; UGT1-05; UGT1-06; UGT1A6S; hUG-BR1; UDPGT 1-6; UGT1A6

Contact

 www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of various lysates using UGT1A6 Rabbit pAb (A17165) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates / proteins: 25 µg per lane.

Blocking buffer: 3 % nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.