# ITPKA Rabbit pAb

www.abclonal.com

ABclonal

Catalog No.: A17503

## **Basic Information**

#### **Observed MW**

55kDa

#### **Calculated MW**

51kDa

## Category

Polyclonal Antibody

## **Applications**

WB, ELISA

## **Cross-Reactivity**

Mouse,Rat

## **Background**

Regulates inositol phosphate metabolism by phosphorylation of second messenger inositol 1,4,5-trisphosphate to Ins(1,3,4,5)P4. The activity of the inositol 1,4,5-trisphosphate 3-kinase is responsible for regulating the levels of a large number of inositol polyphosphates that are important in cellular signaling. Both calcium/calmodulin and protein phosphorylation mechanisms control its activity. It is also a substrate for the cyclic AMP-dependent protein kinase, calcium/calmodulin-dependent protein kinase II, and protein kinase C in vitro.

# **Recommended Dilutions**

**WB** 1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

Gene ID Swiss Prot 3706 P23677

## **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

## **Synonyms**

IP3KA; IP3-3KA; ITPKA

## **Contact**

www.abclonal.com


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

## Validation Data



Western blot analysis of various lysates using ITPKA Rabbit pAb (A17503) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000  $\,$ 

dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 3s.