

[KO Validated] δ-Catenin/p120 Catenin Rabbit pAb

Catalog No.: A18043 KO Validated

Basic Information

Observed MW

100kDa

Calculated MW

108kDa

Category

Polyclonal Antibody

Applications

WB, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

This gene encodes a member of the Armadillo protein family, which function in adhesion between cells and signal transduction. Multiple translation initiation codons and alternative splicing result in many different isoforms being translated. Not all of the full-length natures of the described transcript variants have been determined. Readthrough transcription also exists between this gene and the neighboring upstream thioredoxin-related transmembrane protein 2 (TMX2) gene.

Recommended Dilutions

WB 1:500 - 1:2000

ELISA

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene IDSwiss Prot
1500
060716

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

CAS; p120; BCDS2; CTNND; P120CAS; P120CTN; p120(CAS); p120(CTN); in

Contact

www.abclonal.com


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

Validation Data



Western blot analysis of lysates from wild type (WT) and δ -Catenin/p120 Catenin knockout (KO) HeLa cells, using [KO Validated] δ -Catenin/p120 Catenin Rabbit pAb (A18043) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 5s.