

A19052

Leader in Biomolecular Solutions for Life Science



FGFR3 Rabbit mAb

Catalog No.: A19052

Recombinant

6 Publications

Basic Information

Observed MW

Refer to Figures

Calculated MW

88kDa

Category

SMab Recombinant Monoclonal
Antibody

Applications

IHC-P, ELISA

Cross-Reactivity

Human

CloneNo number

ARC0398

Background

This gene encodes a member of the fibroblast growth factor receptor (FGFR) family, with its amino acid sequence being highly conserved between members and among divergent species. FGFR family members differ from one another in their ligand affinities and tissue distribution. A full-length representative protein would consist of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of the protein interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation. This particular family member binds acidic and basic fibroblast growth hormone and plays a role in bone development and maintenance. Mutations in this gene lead to craniosynostosis and multiple types of skeletal dysplasia.

Recommended Dilutions

IHC-P 1:1000 - 1:5000

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

2261

Swiss Prot

P22607

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

ACH; CEK2; JTK4; CD333; HSFGR3EX; FGFR3

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG


Purification

Affinity purification


Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.




Immunohistochemistry analysis of paraffin-embedded Human pancreas tissue using FGFR3 Rabbit mAb (A19052) at a dilution of 1:2000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human breast cancer tissue using FGFR3 Rabbit mAb (A19052) at a dilution of 1:2000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human testis tissue using FGFR3 Rabbit mAb (A19052) at a dilution of 1:2000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma tissue using FGFR3 Rabbit mAb (A19052) at a dilution of 1:2000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.