ABclonal www.abclonal.com

PADI4 Rabbit pAb

Catalog No.: A1906

Basic Information

Observed MW

74kDa

Calculated MW

74kDa

Category

Polyclonal Antibody

Applications

WB,IF/ICC,ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

This gene is a member of a gene family which encodes enzymes responsible for the conversion of arginine residues to citrulline residues. This gene may play a role in granulocyte and macrophage development leading to inflammation and immune response.

Recommended Dilutions

WB 1:500 - 1:2000

IF/ICC 1:20 - 1:50

ELISA Recommended starting

concentration is 1 µg/mL. Please optimize the concentration based on your specific

assay requirements.

Immunogen Information

Gene ID23569

Swiss Prot
Q9UM07

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

PAD; PAD4; PDI4; PDI5; PADI5; PADI4

Contact

www.abclonal.com


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.
Buffer: PBS with 0.09% Sodium azide,50% glycerol,pH7.3.

Validation Data




Western blot analysis of various lysates using PADI4 Rabbit pAb (A1906) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.



Immunofluorescence analysis of U2OS cells using PADI4 Rabbit pAb (A1906). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.