

A1910

Leader in Biomolecular Solutions for Life Science



Lamin B1 Rabbit pAb

Catalog No.: A1910

57 Publications

Basic Information

Observed MW

68kDa/45kDa

Calculated MW

66kDa

Category

Polyclonal Antibody

Applications

WB,IHC-P,IF/ICC,IP,ChIP,ELISA

Cross-Reactivity

Human,Mouse,Rat

Background

This gene encodes one of the two B-type lamin proteins and is a component of the nuclear lamina. A duplication of this gene is associated with autosomal dominant adult-onset leukodystrophy (ADLD). Alternative splicing results in multiple transcript variants.

Recommended Dilutions

WB 1:500 - 1:1000

IHC-P 1:50 - 1:200

IF/ICC 1:50 - 1:200

IP 0.5µg-4µg antibody for
200µg-400µg extracts
of whole cells

ELISA Recommended starting
concentration is 1
µg/mL. Please optimize
the concentration
based on your specific
assay requirements.

ChIP 5µg antibody for
10µg-15µg of
Chromatin

Immunogen Information

Gene ID

4001

Swiss Prot

P20700

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

LMN; ADLD; LMN2; LMNB; MCPH26; Lamin B1

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Contact



www.abclonal.com

Validation Data



Western blot analysis of various lysates using Lamin B1 Rabbit pAb (A1910) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1min.



Western blot analysis of various lysates using Lamin B1 Rabbit pAb (A1910) at 1:1000 dilution. Jurkat cells were treated with Etoposide (25 uM) at 37°C for 5 hours.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.



Western blot analysis of various lysates using Lamin B1 Rabbit pAb (A1910) at 1:1000 dilution. L929 cells were treated with staurosporine(1 uM) for 3 hour.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 30s.




Immunohistochemistry analysis of paraffin-embedded Human breast cancer using Lamin B1 Rabbit pAb (A1910) at dilution of 1:150 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse kidney using Lamin B1 Rabbit pAb (A1910) at dilution of 1:150 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Rat brain using Lamin B1 Rabbit pAb (A1910) at dilution of 1:150 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunofluorescence analysis of C6 cells using Lamin B1 Rabbit pAb (A1910) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of NIH/3T3 cells using Lamin B1 Rabbit pAb (A1910) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.




Confocal immunofluorescence analysis of HeLa cells using Lamin B1 Rabbit pAb (A1910) at dilution of 1:200. Blue: DAPI for nuclear staining.



Confocal immunofluorescence analysis of HeLa cells using Lamin B1 Rabbit pAb (A1910) at dilution of 1:200. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of PC-12 cells using Lamin B1 Rabbit pAb (A1910) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of U2OS cells using Lamin B1 Rabbit pAb (A1910) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.


Validation Data




Immunoprecipitation analysis of 300 µg extracts of HeLa cells using 3 µg Lamin B1 antibody (A1910). Western blot was performed from the immunoprecipitate using Lamin B1 antibody (A1910) at a dilution of 1:1000.



Immunoprecipitation analysis of 300 µg extracts of HeLa cells using 3 µg Lamin B1 antibody (A1910). Western blot was performed from the immunoprecipitate using Lamin B1 (A1910) at a dilution of 1:1000.




Immunoprecipitation of Lamin B1 in 300 µg extracts from Jurkat cells using 0.5 µg Lamin B1 Rabbit pAb (A1910). Western blot analysis was performed using Lamin B1 Rabbit pAb (A1910) at 1:1000 dilution.



Chromatin immunoprecipitation analysis of extracts of HeLa cells, using Lamin B1 Rabbit pAb antibody (A1910) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.

Validation Data



Chromatin immunoprecipitation was performed with cross-linked chromatin from HeLa, using Lamin-B1 Rabbit mAb (A1910) and rabbit IgG(AC042). The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram compares the ratio of the immunoprecipitated DNA versus the input.