

A19130

Leader in Biomolecular Solutions for Life Science



Transferrin Rabbit mAb

Catalog No.: A19130

Recombinant

7 Publications

Basic Information

Observed MW

77kDa

Calculated MW

77kDa

Category

SMab Recombinant Monoclonal
Antibody

Applications

WB,IHC-P,ELISA

Cross-Reactivity

Human

CloneNo number

ARC0338

Recommended Dilutions

WB 1:1000 - 1:2000

IHC-P 1:100 - 1:1000

ELISA Recommended starting
concentration is 1
µg/mL. Please optimize
the concentration
based on your specific
assay requirements.

Contact

 www.abclonal.com

Background

This gene encodes a glycoprotein with an approximate molecular weight of 76.5 kDa. It is thought to have been created as a result of an ancient gene duplication event that led to generation of homologous C and N-terminal domains each of which binds one ion of ferric iron. The function of this protein is to transport iron from the intestine, reticuloendothelial system, and liver parenchymal cells to all proliferating cells in the body. This protein may also have a physiologic role as granulocyte/pollen-binding protein (GPBP) involved in the removal of certain organic matter and allergens from serum.

Immunogen Information

Gene ID

7018

Swiss Prot

P02787

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 50-150 of human Transferrin (P02787).

Synonyms

TFQTL1; PRO1557; PRO2086; HEL-S-71p; Transferrin

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,0.05% BSA,50% glycerol,pH7.3.

Validation Data



Western blot analysis of lysates from Hep G2 cells using Transferrin Rabbit mAb (A19130) at 1:1000 dilution incubated overnight at 4°C. Hep G2 cells were treated with BFA (30 ng/mL) at 37°C for 1.5 hours.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 30 µg per lane.

Blocking buffer: 3 % nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.