PLA2G2A Rabbit mAb

Catalog No.: A19252 Recombinant

3 Publications



Basic Information

Observed MW 14kDa

Calculated MW 16kDa

Category SMab Recombinant Monoclonal Antibody

Applications WB,ELISA

Cross-Reactivity Human, Mouse, Rat

CloneNo number ARC2416

Recommended Dilutions

1:1000 - 1:6000
Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Background

The protein encoded by this gene is a member of the phospholipase A2 family (PLA2). PLA2s constitute a diverse family of enzymes with respect to sequence, function, localization, and divalent cation requirements. This gene product belongs to group II, which contains secreted form of PLA2, an extracellular enzyme that has a low molecular mass and requires calcium ions for catalysis. It catalyzes the hydrolysis of the sn-2 fatty acid acyl ester bond of phosphoglycerides, releasing free fatty acids and lysophospholipids, and thought to participate in the regulation of the phospholipid metabolism in biomembranes. Several alternatively spliced transcript variants with different 5' UTRs have been found for this gene.

Immunogen Information

Gene ID 5320 Swiss Prot P14555

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 65-144 of human PLA2G2A (P14555).

Synonyms

MOM1; PLA2; PLA2B; PLA2L; PLA2S; PLAS1; sPLA2; PLA2G2A

Contact

Product Information

www.abclonal.com

Source Rabbit **Isotype** IgG **Purification** Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,0.05% BSA,50% glycerol,pH7.3.

Validation Data



Western blot analysis of various lysates using PLA2G2A Rabbit mAb (A19252) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 5s.