# ABclonal® www.abclonal.com

## [KD Validated] RAB27A Rabbit pAb

Catalog No.: A1934 5 Publications

## **Basic Information**

#### **Observed MW**

27kDa

#### **Calculated MW**

25kDa

#### Category

Polyclonal Antibody

## **Applications**

WB,IF/ICC,ELISA

## **Cross-Reactivity**

Human, Mouse

## **Background**

The protein encoded by this gene belongs to the small GTPase superfamily, Rab family. The protein is membrane-bound and may be involved in protein transport and small GTPase mediated signal transduction. Mutations in this gene are associated with Griscelli syndrome type 2. Alternative splicing occurs at this locus and four transcript variants encoding the same protein have been identified.

## **Recommended Dilutions**

**WB** 1:500 - 1:1000

**IF/ICC** 1:50 - 1:200

**ELISA** Recommended starting

concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID**5873

Swiss Prot
P51159

## **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

#### **Synonyms**

GS2; RAM; RAB27; HsT18676; [KD Validated] RAB27A

## **Contact**

www.abclonal.com

## **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

## **Validation Data**




Western blot analysis of lysates from wild type(WT) and RAB27A knockdown (KD) U-87MG cells, using [KD Validated] RAB27A Rabbit pAb (A1934) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.



Immunofluorescence analysis of HepG2 cells using [KD Validated] RAB27A Rabbit pAb (A1934) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.