# ABclonal www.abclonal.com

## EI24 Rabbit pAb

Catalog No.: A19437

## **Basic Information**

#### **Observed MW**

32kDa

#### **Calculated MW**

39kDa

## **Category**

Polyclonal Antibody

#### **Applications**

WB, ELISA

## **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

This gene encodes a putative tumor suppressor and has higher expression in p53-expressing cells than in control cells and is an immediate-early induction target of p53-mediated apoptosis. The encoded protein may suppress cell growth by inducing apoptotic cell death through the caspase 9 and mitochondrial pathways. This gene is located on human chromosome 11q24, a region frequently altered in cancers. Alternative splicing results in multiple transcript variants. Pseudogenes of this gene have been defined on chromosomes 1, 3, 7, and 8.

## **Recommended Dilutions**

WB 1:1000 - 1:5000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID**9538

Swiss Prot
014681

## **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

## **Synonyms**

EPG4; PIG8; TP53I8; EI24

## Contact

www.abclonal.com

## **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

## Validation Data



Western blot analysis of various lysates, using EI24 Rabbit pAb (A19437) at 1:2000

dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000  $\,$ 

dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.