ABclonal www.abclonal.com

CTIP2/BCL11B Rabbit mAb

Catalog No.: A19804 Recombinant 1 Publications

Basic Information

Observed MW

120-130kDa

Calculated MW

96kDa

Category

SMab Recombinant Monoclonal Antibody

Applications

WB,IHC-P,ELISA

Cross-Reactivity

Human, Mouse

CloneNo number

ARC2329

Background

This gene encodes a C2H2-type zinc finger protein and is closely related to BCL11A, a gene whose translocation may be associated with B-cell malignancies. Although the specific function of this gene has not been determined, the encoded protein is known to be a transcriptional repressor, and is regulated by the NURD nucleosome remodeling and histone deacetylase complex. Four alternatively spliced transcript variants encoding distinct isoforms have been found for this gene.

Recommended Dilutions

WB 1:1000 - 1:6000

IHC-P 1:50 - 1:200

Recommended starting **ELISA**

> concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

64919 **Immunogen**

Gene ID

A synthetic peptide corresponding to a sequence within amino acids 450-550 of human CTIP2/BCL11B (Q9C0K0).

Swiss Prot

Q9C0K0

Synonyms

ATL1; RIT1; CTIP2; IMD49; CTIP-2; IDDFSTA; SMARCM2; ZNF856B; ATL1-beta; ATL1alpha; ATL1-delta; ATL1-gamma; hRIT1-alpha; CTIP2/BCL11B

Contact

€ www.abclonal.com

Product Information

Immunogen Information


Source **Isotype Purification** Rabbit Affinity purification IgG

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of various lysates using CTIP2/BCL11B Rabbit mAb (A19804) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000


dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.



Immunohistochemistry analysis of paraffin-embedded Mouse brain using CTIP2/BCL11B Rabbit mAb (A19804) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.