

A19815

Leader in Biomolecular Solutions for Life Science



# Histone H4 Rabbit mAb

Catalog No.: A19815

Recombinant

1 Publications

## Basic Information

### Observed MW

11kDa

### Calculated MW

11kDa

### Category

SMab Recombinant Monoclonal Antibody

### Applications

WB,IHC-P,ELISA

### Cross-Reactivity

Human,Mouse,Rat,Other (Wide Range Predicted)

### CloneNo number

ARC2340

## Recommended Dilutions

**WB** 1:1000 - 1:6000

**IHC-P** 1:100 - 1:1000

**ELISA** Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## Contact



[www.abclonal.com](http://www.abclonal.com)

## Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H4 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element.

## Immunogen Information

**Gene ID**  
8359

**Swiss Prot**  
P62805

### Immunogen

A synthetic peptide corresponding to a sequence within amino acids 24-103 of human Histone H4 (P62805).

### Synonyms

H4/p; H4C1; H4C2; H4C3; H4C4; H4C5; H4C6; H4C8; H4C9; H4-16; H4C11; H4C12; H4C13; H4C14; H4C15; HIST4H4; Histone H4

## Product Information

**Source**  
Rabbit

**Isotype**  
IgG


**Purification**  
Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,0.05% BSA,50% glycerol,pH7.3.


## Validation Data




Western blot analysis of various lysates using Histone H4 Rabbit mAb (A19815) at 1:1000 dilution.  
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Enhanced Kit (RM00021).  
Exposure time: 10s.



Immunohistochemistry analysis of paraffin-embedded Rat liver using Histone H4 Rabbit mAb (A19815) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human colon using Histone H4 Rabbit mAb (A19815) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse kidney using Histone H4 Rabbit mAb (A19815) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.