

SARS-CoV-2 N Protein Rabbit mAb, BSA and glycerol free

Catalog No.: A20021 Recombinant 17 Publications

Basic Information

Observed MW

46kDa (Recombinant SARS-CoV-2 Nucleocapsid Protein)

Calculated MW

46kDa

Category

SMab Recombinant Monoclonal Antibody

Applications

WB,ELISA

Cross-Reactivity

SARS-CoV-2

CloneNo number

ARC5077-02

Background

Packages the positive strand viral genome RNA into a helical ribonucleocapsid (RNP and plays a fundamental role during virion assembly through its interactions with the viral genome and membrane protein M. Plays an important role in enhancing the efficiency of subgenomic viral RNA transcription as well as viral replication. May modulate transforming growth factor-beta signaling by binding host SMAD3.

Recommended Dilutions

WB 1:1000 - 1:6000

ELISA

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID43740575

Swiss Prot
P59595

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

Nucleoprotein; NP; SARS-CoV-2 N Protein

Contact

www.abclonal.com

Product Information


SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C Avoid freeze / thaw cycles.

Buffer: PBS, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of Recombinant SARS-CoV-2 Nucleocapsid Protein (RP01264LQ), using SARS-CoV-2 N Protein Rabbit mAb,BSA and glycerol free (A20021) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 30 ng/10 ng/5 ng/2 ng per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 5s.