# ABclonal®

# **CDC45 Rabbit pAb**

Catalog No.: A2047

# **Basic Information**

#### **Observed MW**

60kDa

#### **Calculated MW**

66kDa

#### Category

Polyclonal Antibody

#### **Applications**

WB, ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

The protein encoded by this gene was identified by its strong similarity with Saccharomyces cerevisiae Cdc45, an essential protein required to the initiation of DNA replication. Cdc45 is a member of the highly conserved multiprotein complex including Cdc6/Cdc18, the minichromosome maintenance proteins (MCMs) and DNA polymerase, which is important for early steps of DNA replication in eukaryotes. This protein has been shown to interact with MCM7 and DNA polymerase alpha. Studies of the similar gene in Xenopus suggested that this protein play a pivotal role in the loading of DNA polymerase alpha onto chromatin. Alternate splicing results in multiple transcript variants.

#### **Recommended Dilutions**

WB 1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

 Gene ID
 Swiss Prot

 8318
 075419

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

#### **Synonyms**

CDC45L; MGORS7; CDC45L2; PORC-PI-1; CDC45

### **Contact**

www.abclonal.com


#### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# Validation Data



Western blot analysis of various lysates using CDC45 Rabbit pAb (A2047) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000  $\,$ 

dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 15s.