

A20683

Leader in Biomolecular Solutions for Life Science



UGT2B17 Rabbit pAb

Catalog No.: A20683

Basic Information

Observed MW

Refer to figures

Calculated MW

61kDa

Category

Polyclonal Antibody

Applications

IF/ICC,ELISA

Cross-Reactivity

Human

Background

This gene encodes a member of the uridine diphosphoglucuronosyltransferase protein family. The encoded enzyme catalyzes the transfer of glucuronic acid from uridine diphosphoglucuronic acid to a diverse array of substrates including steroid hormones and lipid-soluble drugs. This process, known as glucuronidation, is an intermediate step in the metabolism of steroids. Copy number variation in this gene is associated with susceptibility to osteoporosis.

Recommended Dilutions

IF/ICC 1:50 - 1:200

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

7367

Swiss Prot

O75795

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

BMND12; UDPGT2B17; UGT2B17

Contact

 www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

Validation Data



Immunofluorescence analysis of HepG2 cells using UGT2B17 Rabbit pAb (A20683) at dilution of 1:200 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of PC-3 cells using UGT2B17 Rabbit pAb (A20683) at dilution of 1:200 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.