

A20735

Leader in Biomolecular Solutions for Life Science



# Acetyl-Histone H3-K18 Rabbit mAb

Catalog No.: A20735

Recombinant

## Basic Information

### Observed MW

17kDa

### Calculated MW

15kDa

### Category

SMab Recombinant Monoclonal  
Antibody

### Applications

WB,IHC-P,ChIP,ChIP-  
seq,ELISA,DB,CUT&Tag

### Cross-Reactivity

Human,Mouse,Rat,Other (Wide Range  
Predicted)

### CloneNo number

ARC53056

## Recommended Dilutions

**WB** 1:500 - 1:1000

**DB** 1:10000 - 1:100000

**IHC-P** 1:50 - 1:200

**ELISA** Recommended starting  
concentration is 1  
µg/mL. Please optimize  
the concentration  
based on your specific  
assay requirements.

**ChIP** 5µg antibody for  
5µg-10µg of Chromatin

**ChIP-seq** 1:50 - 1:200

**CUT&Tag** 10<sup>5</sup> cells /1 µg

## Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3.

## Immunogen Information

### Gene ID

8290/8350

### Swiss Prot

Q16695/P68431

### Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

### Synonyms

H3/A; H3C2; H3C3; H3C4; H3C6; H3C7; H3C8; H3FA; H3C10; H3C11; H3C12; HIST1H3A; H3C1; Acetyl-Histone H3-K18

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification

Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.

## Contact




[www.abclonal.com](http://www.abclonal.com)


Validation Data




Western blot analysis of various lysates using Acetyl-Histone H3-K18 Rabbit mAb (A20680) at 1:1000 dilution. HeLa cells and NIH/3T3 cells and C6 cells were treated with TSA (1  $\mu$ M) at 37°C for 18 hours.  
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25 $\mu$ g per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 180s.




Dot-blot analysis of all sorts of peptides using Acetyl-Histone H3-K18 Rabbit mAb (A20735) at 1:100000 dilution.




Immunohistochemistry analysis of paraffin-embedded Human cervix cancer tissue using Acetyl-Histone H3-K18 Rabbit mAb (A20735) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma tissue using Acetyl-Histone H3-K18 Rabbit mAb (A20735) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human colon tissue using Acetyl-Histone H3-K18 Rabbit mAb (A20735) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse brain tissue using Acetyl-Histone H3-K18 Rabbit mAb (A20735) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse intestine tissue using Acetyl-Histone H3-K18 Rabbit mAb (A20735) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse kidney tissue using Acetyl-Histone H3-K18 Rabbit mAb (A20735) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse liver tissue using Acetyl-Histone H3-K18 Rabbit mAb (A20735) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse lung tissue using Acetyl-Histone H3-K18 Rabbit mAb (A20735) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse spleen tissue using Acetyl-Histone H3-K18 Rabbit mAb (A20735) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Rat brain tissue using Acetyl-Histone H3-K18 Rabbit mAb (A20735) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat colon tissue using Acetyl-Histone H3-K18 Rabbit mAb (A20735) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat liver tissue using Acetyl-Histone H3-K18 Rabbit mAb (A20735) at a dilution of 1:200 (40x lens). High pressure antigen




Immunohistochemistry analysis of paraffin-embedded Rat lung tissue using Acetyl-Histone H3-K18 Rabbit mAb (A20735) at a dilution of 1:200 (40x lens). High pressure antigen


## Validation Data

retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.


retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Chromatin immunoprecipitation analysis of extracts of HeLa cells, using Acetyl-Histone H3-K18 Rabbit mAb (A20735) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.



Chromatin immunoprecipitations were performed with cross-linked chromatin from 293T cells and Acetyl-Histone H3-K18 Rabbit mAb (A20735). The ChIP sequencing results indicate the enrichment pattern of Acetyl-Histone H3-K18 in selected genomic region and representative gene loci (RPL30), as shown in figure.



CUT&Tag was performed using the CUT&Tag Assay Kit (pAG-Tn5) for Illumina(RK20265) from 10<sup>5</sup> K562 cells with 1 µg Acetyl-Histone H3-K18 Rabbit mAb (A20735), along with a Goat Anti-Rabbit IgG(H+L). The CUT&Tag results indicate the enrichment pattern of H3K18me1 in representative gene loci (RPL30), as shown in figure.