

A21980

Leader in Biomolecular Solutions for Life Science



POLR2A (N-terminal) Rabbit mAb

Catalog No.: A21980

Recombinant

1 Publications

Basic Information

Observed MW

250kDa

Calculated MW

217kDa

Category

SMab Recombinant Monoclonal
Antibody

Applications

WB,IP,ChIP,ELISA

Cross-Reactivity

Human,Mouse,Rat

CloneNo number

ARC54153

Background

This gene encodes the largest subunit of RNA polymerase II, the polymerase responsible for synthesizing messenger RNA in eukaryotes. The product of this gene contains a carboxy terminal domain composed of heptapeptide repeats that are essential for polymerase activity. These repeats contain serine and threonine residues that are phosphorylated in actively transcribing RNA polymerase. In addition, this subunit, in combination with several other polymerase subunits, forms the DNA binding domain of the polymerase, a groove in which the DNA template is transcribed into RNA.

Recommended Dilutions

WB 1:2000 - 1:8000

IP 0.5µg-4µg antibody for
400µg-600µg extracts
of whole cells

ChIP 5µg antibody for
10µg-15µg of
Chromatin

ELISA Recommended starting
concentration is 1
µg/mL. Please optimize
the concentration
based on your specific
assay requirements.

Immunogen Information

Gene ID

5430

Swiss Prot

P24928

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

RPB1; RPO2; POLR2; POLRA; RPBh1; RPOL2; NEDHIB; RpIILS; hsRPB1; hRPB220; POLR2A (N-terminal)

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Contact




www.abclonal.com


Validation Data




Western blot analysis of lysates from Mouse thymus, using POLR2A (N-terminal) Rabbit mAb (A21980) at 1:2000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 10s.



Western blot analysis of lysates from C6 cells using POLR2A (N-terminal) Rabbit mAb (A21980) at 1:2000 dilution incubated overnight at 4°C.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 45s.



Immunoprecipitation analysis of 600 µg extracts of 293F cells using 3 µg POLR2A (N-terminal) Rabbit mAb (A21980). Western blot was performed from the immunoprecipitate using POLR2A (N-terminal) Rabbit mAb (A21980) at a dilution of 1:1000.



Chromatin immunoprecipitation analysis of extracts of 293F cells, using POLR2A (N-terminal) Rabbit mAb (A21980) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.