ABclonal®

AIFM2/FSP1 Rabbit mAb

Catalog No.: A22278 Recombinant 3 Publications

Basic Information

Observed MW

44kDa

Calculated MW

41kDa

Category

SMab Recombinant Monoclonal Antibody

Applications

WB,IHC-P,ELISA

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC56793

Background

This gene encodes a flavoprotein oxidoreductase that binds single stranded DNA and is thought to contribute to apoptosis in the presence of bacterial and viral DNA. The expression of this gene is also found to be induced by tumor suppressor protein p53 in colon cancer cells.

Recommended Dilutions

WB 1:10000 - 1:40000

IHC-P 1:200 - 1:800

ELISA Recommended starting

concentration is 1 µg/mL. Please optimize the concentration based on your specific

assay requirements.

Immunogen Information

Gene IDSwiss Prot
84883
Q9BRQ8

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 43-371 of human AIFM2/FSP1 (NP_116186.1).

Synonyms

AMID; FSP1; PRG3; AIFM2/FSP1

Contact

www.abclonal.com


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide, 0.05% BSA, 50% glycerol, pH7.3.



Western blot analysis of various lysates, using AIFM2/FSP1 Rabbit mAb (A22278) at1:20000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000


dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.



Immunohistochemistry analysis of paraffin-embedded Mouse liver tissue using AIFM2/FSP1 Rabbit mAb (A22278) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat liver tissue using AIFM2/FSP1 Rabbit mAb (A22278) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.