# ABclonal® www.abclonal.com

## KDM3A Rabbit pAb

Catalog No.: A2322 3 Publications

## **Basic Information**

#### **Observed MW**

147kDa

#### **Calculated MW**

147kDa

#### Category

Polyclonal Antibody

### **Applications**

WB, ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

Enables androgen receptor binding activity; histone H3-methyl-lysine-9 demethylase activity; and iron ion binding activity. Involved in several processes, including androgen receptor signaling pathway; formaldehyde biosynthetic process; and histone H3-K9 demethylation. Located in nucleoplasm. Implicated in cervical cancer and colon cancer. Biomarker of Ewing sarcoma; hepatocellular carcinoma; nasopharynx carcinoma; and prostate cancer.

## **Recommended Dilutions**

WB 1:500 - 1:1000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

 Gene ID
 Swiss Prot

 55818
 Q9Y4C1

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

## **Synonyms**

TSGA; JMJD1; JHDM2A; JHMD2A; JMJD1A; KDM3A

## **Contact**

www.abclonal.com


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of lysates from 293T cells, using KDM3A Rabbit pAb (A2322) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.



Western blot analysis of lysates from Mouse testis, using KDM3A Rabbit pAb (A2322) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.