# IL4 Rabbit mAb

Catalog No.: A23277 Recombinant 2 Publications



### **Basic Information**

### **Observed MW**

20-23KDa

### **Calculated MW**

17kDa

#### Category

SMab Recombinant Monoclonal Antibody

### **Applications**

WB, ELISA

### **Cross-Reactivity**

Human

### CloneNo number

ARC52902

### **Recommended Dilutions**

WB 1:1000 - 1:5000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

# <u>Background</u>

The protein encoded by this gene is a pleiotropic cytokine produced by activated T cells. This cytokine is a ligand for interleukin 4 receptor. The interleukin 4 receptor also binds to IL13, which may contribute to many overlapping functions of this cytokine and IL13. STAT6, a signal transducer and activator of transcription, has been shown to play a central role in mediating the immune regulatory signal of this cytokine. This gene, IL3, IL5, IL13, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL13. This gene, IL13 and IL5 are found to be regulated coordinately by several long-range regulatory elements in an over 120 kilobase range on the chromosome. IL4 is considered an important cytokine for tissue repair, counterbalancing the effects of proinflammatory type 1 cytokines, however, it also promotes allergic airway inflammation. Moreover, IL-4, a type 2 cytokine, mediates and regulates a variety of human host responses such as allergic, anti-parasitic, wound healing, and acute inflammation. This cytokine has been reported to promote resolution of neutrophil-mediated acute lung injury. In an allergic response, IL-4 has an essential role in the production of allergen-specific immunoglobin (Ig) E. This proinflammatory cytokine has been observed to be increased in COVID-19 (Coronavirus disease 2019) patients, but is not necessarily associated with severe COVID-19 pathology. Two alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported.

## **Immunogen Information**

**Gene ID**Swiss Prot
3565
P05112

### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### **Synonyms**

BSF1; IL-4; BCGF1; BSF-1; BCGF-1; IL4

### **Contact**

www.abclonal.com

### **Product Information**


SourceIsotypePurificationRabbitIqGAffinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

## Validation Data



Western blot analysis of lysates from Recombinant Human IL-4 Protein (RP00995) using IL4 Rabbit mAb (A23277) at 1:1000 dilution incubated overnight at  $4^{\circ}$ C.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000

dilution.

Lysates/proteins: 10ng,5ng per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.