

A23289

Leader in Biomolecular Solutions for Life Science



## VDR Rabbit mAb

Catalog No.: A23289

Recombinant

1 Publications

### Basic Information

**Observed MW**

48kDa/54kDa

**Calculated MW**

48kDa

**Category**SMab Recombinant Monoclonal  
Antibody**Applications**

WB,IHC-P,IF/ICC,IP,ChIP,ELISA

**Cross-Reactivity**

Human,Mouse,Rat

**CloneNo number**

ARC60266

### Background

This gene encodes vitamin D3 receptor, which is a member of the nuclear hormone receptor superfamily of ligand-inducible transcription factors. This receptor also functions as a receptor for the secondary bile acid, lithocholic acid. Downstream targets of vitamin D3 receptor are principally involved in mineral metabolism, though this receptor regulates a variety of other metabolic pathways, such as those involved in immune response and cancer. Mutations in this gene are associated with type II vitamin D-resistant rickets. A single nucleotide polymorphism in the initiation codon results in an alternate translation start site three codons downstream. Alternatively spliced transcript variants encoding different isoforms have been described for this gene. A recent study provided evidence for translational readthrough in this gene, and expression of an additional C-terminally extended isoform via the use of an alternative in-frame translation termination codon.

### Recommended Dilutions

**WB** 1:500 - 1:1000**IHC-P** 1:500 - 1:2000**IF/ICC** 1:100 - 1:400**IP** 0.5µg-4µg antibody for  
200µg-400µg extracts  
of whole cells**ELISA** Recommended starting  
concentration is 1  
µg/mL. Please optimize  
the concentration  
based on your specific  
assay requirements.**ChIP** 5µg antibody for  
5µg-10µg of Chromatin

### Immunogen Information

**Gene ID**

7421

**Swiss Prot**

P11473

**Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

**Synonyms**

NR1I1; PPP1R163; VDR

### Product Information

**Source**

Rabbit

**Isotype**

IgG

**Purification**

Affinity purification

**Storage**


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

### Contact

[www.abclonal.com](http://www.abclonal.com)

## Validation Data



Western blot analysis of various lysates, using VDR Rabbit mAb (A23289) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 180s.




Immunohistochemistry analysis of paraffin-embedded Human cervix cancer tissue using VDR Rabbit mAb (A23289) at a dilution of 1:500 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse colon tissue using VDR Rabbit mAb (A23289) at a dilution of 1:500 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human colon tissue using VDR Rabbit mAb (A23289) at a dilution of 1:500 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma tissue using VDR Rabbit mAb (A23289) at a dilution of 1:500 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Rat kidney tissue using VDR Rabbit mAb (A23289) at a dilution of 1:500 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Rat liver tissue using VDR Rabbit mAb (A23289) at a dilution of 1:500 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.


Validation Data



Immunofluorescence analysis of paraffin-embedded rat kidney using VDR Rabbit mAb (A23289) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of T-47D cells using VDR Rabbit mAb (A23289) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunoprecipitation analysis of 300 µg extracts of Jurkat cells using 3 µg VDR Rabbit mAb (A23289). Western blot was performed from the immunoprecipitate using VDR Rabbit mAb (A23289) at a dilution of 1:1000.