# eEF1A1 Rabbit mAb

Catalog No.: A23515 Recombinant 3 Publications



## **Basic Information**

## **Observed MW**

50kDa

#### **Calculated MW**

50kDa

### **Category**

SMab Recombinant Monoclonal Antibody

### **Applications**

WB,IHC-P,IF/ICC,ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

#### CloneNo number

ARC55875

## **Background**

This gene encodes an isoform of the alpha subunit of the elongation factor-1 complex, which is responsible for the enzymatic delivery of aminoacyl tRNAs to the ribosome. This isoform (alpha 1) is expressed in brain, placenta, lung, liver, kidney, and pancreas, and the other isoform (alpha 2) is expressed in brain, heart and skeletal muscle. This isoform is identified as an autoantigen in 66% of patients with Felty syndrome. This gene has been found to have multiple copies on many chromosomes, some of which, if not all, represent different pseudogenes.

### **Recommended Dilutions**

WB 1:1000 - 1:5000
IHC-P 1:50 - 1:200
IF/ICC 1:50 - 1:200

**ELISA** Recommended starting

concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

### **Contact**

www.abclonal.com

## **Immunogen Information**

Gene ID	Swiss Prot
1915	P68104

## Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### **Synonyms**

CCS3; EF1A; PTI1; CCS-3; EE1A1; EEF-1; EEF1A; EF-Tu; EF1A1; LENG7; eEF1A-1; GRAF-1EF; EF1alpha1; eEF1A1


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide, 0.05% BSA, 50% glycerol, pH7.3.



Western blot analysis of various lysates, using eEF1A1 Rabbit mAb (A23515) at 1:2000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 0.2s.




Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma tissue using eEF1A1 Rabbit mAb (A23515) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse testis tissue using eEF1A1 Rabbit mAb (A23515) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human lung adenocarcinoma tissue using eEF1A1 Rabbit mAb (A23515) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat brain tissue using eEF1A1 Rabbit mAb (A23515) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse kidney tissue using eEF1A1 Rabbit mAb (A23515) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Rat kidney tissue using eEF1A1 Rabbit mAb (A23515) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.

## **Validation Data**



Immunohistochemistry analysis of paraffin-embedded Rat spleen tissue using eEF1A1 Rabbit mAb (A23515) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunofluorescence analysis of C6 cells using eEF1A1 Rabbit mAb (A23515) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.