

# Fibroblast activation protein-α (FAP) Rabbit mAb

Catalog No.: A23789 Recombinant 4 Publications

## **Basic Information**

#### **Observed MW**

95kDa

### **Calculated MW**

26kDa/87kDa

### Category

SMab Recombinant Monoclonal Antibody

### **Applications**

WB,IF/ICC,ELISA

# **Cross-Reactivity**

Human, Mouse, Rat

### CloneNo number

ARC61700

# **Background**

The protein encoded by this gene is a homodimeric integral membrane gelatinase belonging to the serine protease family. It is selectively expressed in reactive stromal fibroblasts of epithelial cancers, granulation tissue of healing wounds, and malignant cells of bone and soft tissue sarcomas. This protein is thought to be involved in the control of fibroblast growth or epithelial-mesenchymal interactions during development, tissue repair, and epithelial carcinogenesis. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

# **Recommended Dilutions**

**WB** 1:1000 - 1:4000

**IF/ICC** 1:200 - 1:800

**ELISA** Recommended starting

concentration is 1 µg/mL. Please optimize the concentration based on your specific

assay requirements.

# **Immunogen Information**

**Gene ID**2191

Swiss Prot
Q12884

### **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

### **Synonyms**

FAP; DPPIV; FAPA; FAPalpha; SIMP; prolyl endopeptidase FAP; Fibroblast activation protein- $\alpha$  (FAP)

# Contact

www.abclonal.com

### **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

# **Validation Data**



Western blot analysis of various lysates, using Fibroblast activation protein- $\alpha$  (FAP) Rabbit mAb (A23789) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.


Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.





Immunofluorescence analysis of NIH/3T3 cells using Fibroblast activation protein- $\alpha$  (FAP) Rabbit mAb (A23789) at dilution of 1:200 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.

Immunofluorescence analysis of U-87MG cells using Fibroblast activation protein- $\alpha$  (FAP) Rabbit mAb (A23789) at dilution of 1:200 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.