# ABclomal www.abclonal.com

## γ-Tubulin Rabbit mAb

Catalog No.: A23935 Recombinant

## **Basic Information**

#### **Observed MW**

50kDa

#### **Calculated MW**

51kDa

## **Category**

SMab Recombinant Monoclonal Antibody

## **Applications**

WB, ELISA

## **Cross-Reactivity**

Human, Mouse, Rat

#### CloneNo number

ARC54998

## **Background**

This gene encodes a member of the tubulin superfamily. The encoded protein localizes to the centrosome where it binds to microtubules as part of a complex referred to as the gamma-tubulin ring complex. The protein mediates microtubule nucleation and is required for microtubule formation and progression of the cell cycle. A pseudogene of this gene is found on chromosome 7.

## **Recommended Dilutions**

**WB** 1:500 - 1:1000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID**7283

Swiss Prot
P23258

## **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### **Synonyms**

TUBG; GCP-1; CDCBM4; TUBGCP1; γ-Tubulin

## **Contact**

www.abclonal.com

## **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

## **Validation Data**



Western blot analysis of various lysates, using  $\gamma$ -Tubulin Rabbit mAb (A23935) at1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 20s.