

A2461

Leader in Biomolecular Solutions for Life Science



PIK3CG Rabbit pAb

Catalog No.: A2461

Basic Information

Observed MW

112kDa

Calculated MW

126kDa

Category

Polyclonal Antibody

Applications

WB,IP

Cross-Reactivity

Human,Mouse

Background

Phosphoinositide 3-kinases (PI3Ks) phosphorylate inositol lipids and are involved in the immune response. The protein encoded by this gene is a class I catalytic subunit of PI3K. Like other class I catalytic subunits (p110-alpha p110-beta, and p110-delta), the encoded protein binds a p85 regulatory subunit to form PI3K. This gene is located in a commonly deleted segment of chromosome 7 previously identified in myeloid leukemias. Several transcript variants encoding the same protein have been found for this gene.

Recommended Dilutions

WB 1:500 - 1:1000

IP 1:20 - 1:100

Immunogen Information

Gene ID

5294

Swiss Prot

P48736

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

PI3K; PIK3; IMD97; PI3CG; PI3Kgamma; p110gamma; p120-PI3K; PIK3CG

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Validation Data



Western blot analysis of lysates from Jurkat cells, using PIK3CG Rabbit pAb (A2461).
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.